Animal Specialty and Emergency Hospital 5775 Schenck Avenue ~ Rockledge, FL 32955 T: (321) 752-7600 ~ F: (321) 752-4882

Potential chemotherapy side effects

The following is a list and explanation of potential chemotherapy side effects.

Vomiting, nausea, and diarrhea – Gastrointestinal upset is a common side effect of most chemotherapy drugs, usually noticed 24–48 hours and/or 3–5 days post-treatment.

Vomiting: Remove food and only allow small amounts of water for 12 hours. Then, introduce a bland diet slowly over the next 12 hours. Please call the hospital if vomit contains blood, is severe (3 times or more per hour) or does not resolve within 24 hours. Prescription anti-nausea medication may be required.

Diarrhea: Diarrhea often resolves without treatment. If diarrhea is mild to moderate, feed your pet his/her normal diet and observe closely for other symptoms (vomiting, fever, lethargy, etc.). Please call the hospital if diarrhea is severe, contains blood, is accompanied by other symptoms, or lasts longer than 24 hours.

Injection site irritation – Most chemotherapy drugs are administered intravenously, and we take every precaution to prevent extravasation (injection of the drug outside of the vein). Some chemotherapy drugs can cause severe skin irritation, necrosis, and sloughing if extravasation occurs. Most injection site irritation will begin to show within 24-48 hours. Please call the hospital if you notice any irritation or discomfort at the site of an injection.

Things to look for:

Licking or chewing at injection site Redness or rash that resembles a burn Pain or sensitivity in the area that the injection was given

<u>Fever</u> – A fever can be caused by stress, or infection due to a compromised immune system. You should designate a digital rectal thermometer for your pet. If your pet feels warm to the touch or does not seem to feel well you should take his/her temperature. Your pet's temperature should be somewhere between 100.0 and 102.0. Anytime your pet's temperature is 103.0 or higher you should contact the hospital immediately, as this could indicate life threatening infection in a chemotherapy patient.

Things to look for:

Common symptoms in a pet with a fever are lethargy, depression and inappetance.

<u>Allergic reaction</u> – Some chemotherapy drugs can cause an allergic reaction. These allergic reactions can vary from mild to life threatening. Prior to giving these drugs we will administer an antihistamine and will check your pet for signs of allergic reaction before you take him/her home. Most allergic reactions will occur within 20 minutes of receiving chemotherapy, but you should observe your pet closely for 24 hours.

Things to look for:

Hives or rash on your pet's abdomen, ears, feet or gums. Difficulty breathing, facial swelling, vomiting, excessive salivation

<u>Petechia</u> – Petechia or bruising is usually caused by a low platelet count and could indicate bone marrow suppression. Petechia will usually appear as reddish or purple spots varying in size and are usually most visible on the abdomen. If you notice petechia or any abnormal bleeding please contact the hospital immediately.

Things to look for: Bleeding from your pet's mouth or nose Blood in you pet's urine or feces. *Please note that hemorrhagic cystitis can occur from certain chemotherapy drugs without an underlying platelet problem.* Petechia anywhere on your pet's body

<u>Alopecia and hyperpigmentation</u> – Alopecia and hyperpigmentation are not harmful to your pet. Alopecia (hair loss) is most common in breeds whose hair grows continuously (Poodles, Terriers, Old English Sheepdogs, etc.). Alopecia may involve the entire body, or only focal areas such as the abdomen or whiskers. Hyperpigmentation (darkened skin color) can occur in any breed. Alopecia often resolves, but hyperpigmentation is usually permanent.

Hemorrhagic cystitis – The chemotherapy drug cyclophosphamide can irritate the inner lining of the urinary bladder, causing it to bleed. This condition is known as hemorrhagic cystitis. It is important that your pet have free access to fresh water and be taken outside to urinate frequently. This can be a serious condition, so contact the hospital if you notice blood in your pet's urine.

Please feel free to contact us if you have any questions or concerns. Any changes in your pet's attitude or activity level should be noted and observed. If you pet does not feel well or is just 'not acting right' please let us know as this could be an early indication of a more serious problem.